

Tilburg University

A representation theorem for $(\text{trAp})_{1/p}$

Magnus, J.R.

Published in:
Linear Algebra and its Applications

Publication date:
1987

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):
Magnus, J. R. (1987). A representation theorem for $(\text{trAp})_{1/p}$. *Linear Algebra and its Applications*, 95, 127-134.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

A Representation Theorem for $(\text{tr } A^p)^{1/p}$

Jan R. Magnus

London School of Economics

Houghton Street

London, WC2A 2AE England

Submitted by Ingram Olkin

ABSTRACT

An inequality for positive semidefinite matrices is proved, and from it a quasilinear representation of $(\text{tr } A^p)^{1/p}$ is obtained. From this representation follow matrix versions of the inequalities of Hölder and Minkowski.

1. INTRODUCTION

The fundamental inequalities of Hölder and Minkowski can be proved in a variety of ways; see Hardy, Littlewood, and Pólya (1952). A particularly interesting proof of both inequalities is based on the following lemma, of interest in itself, which is easily established by the method of Lagrange or otherwise.

LEMMA 1.¹ *Let $p > 1$, $q = p/(p - 1)$, and $a_i \geq 0$ ($i = 1, \dots, n$). Then*

$$\sum_{i=1}^n a_i x_i \leq \left(\sum_{i=1}^n a_i^p \right)^{1/p} \quad (1.1)$$

for every set of nonnegative quantities x_1, x_2, \dots, x_n satisfying $\sum_{i=1}^n x_i^q = 1$. Equality in (1.1) occurs if and only if $a_1 = a_2 = \dots = a_n = 0$ or $x_i^q = a_i^p / \sum_{j=1}^n a_j^p$ ($i = 1, \dots, n$).

¹See Beckenbach and Bellman (1961, Theorem 5, p. 23).

An immediate and easy consequence of Lemma 1 is

HÖLDER'S INEQUALITY. *If $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$ are vectors in the nonnegative orthant of \mathbb{R}^n and $0 < \alpha < 1$, then*

$$\sum_{i=1}^n x_i^\alpha y_i^{1-\alpha} \leq \left(\sum_{i=1}^n x_i \right)^\alpha \left(\sum_{i=1}^n y_i \right)^{1-\alpha} \quad (1.2)$$

with equality if and only if x and y are linearly dependent.

If we formalate the inequality (1.1) as a maximization problem, we get

$$\max_R \sum_{i=1}^n a_i x_i = \left(\sum_{i=1}^n a_i^p \right)^{1/p}, \quad (1.3)$$

where R is the region defined by

$$\sum_{i=1}^n x_i^q = 1, \quad x_i \geq 0.$$

The idea of expressing a *nonlinear* function, such as $(\sum a_i^p)^{1/p}$, as an envelope of *linear* functions goes back to Minkowski and was used extensively by Bellman and others. This technique is called *quasilinearization*. A direct consequence of the quasilinear representation (1.3) is

MINKOWSKI'S INEQUALITY. *If $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$ are vectors in the nonnegative orthant of \mathbb{R}^n and $p > 1$, then*

$$\left(\sum_{i=1}^n (x_i + y_i)^p \right)^{1/p} \leq \left(\sum_{i=1}^n x_i^p \right)^{1/p} + \left(\sum_{i=1}^n y_i^p \right)^{1/p} \quad (1.4)$$

with equality if and only if x and y are linearly dependent.

The purpose of this paper is to extend the inequality (1.1) to positive semidefinite matrices, thus obtaining a quasilinear representation of $(\text{tr } A^p)^{1/p}$. This is achieved in Theorem 5. Interesting matrix versions of the inequalities of Hölder and Minkowski are then easily derived from this representation (Theorems 6 and 7). Several preliminary results, of interest in themselves, are also reported.

2. A NECESSARY AND SUFFICIENT CONDITION FOR DIAGONALITY

Let us begin by stating the following simple but useful result concerning the conditions under which a real symmetric (or positive definite) matrix is diagonal.

THEOREM 1. *A positive definite matrix is diagonal if and only if the product of its diagonal elements is equal to its determinant. A real symmetric matrix is diagonal if and only if its diagonal elements and its eigenvalues coincide.*

Proof. A proof by induction and perturbation is straightforward and therefore omitted. A variety of other proofs can also be constructed. ■

3. KARAMATA'S INEQUALITY

Let $x = (x_1, x_2, \dots, x_n)$ and $y = (y_1, y_2, \dots, y_n)$ be two vectors in \mathbb{R}^n . We shall say that x is *majorized* by y , and write

$$(x_1, \dots, x_n) \prec (y_1, \dots, y_n),$$

when the following three relations are satisfied:

$$x_1 + x_2 + \dots + x_n = y_1 + y_2 + \dots + y_n;$$

$$x_1 \geq x_2 \geq \dots \geq x_n, \quad y_1 \geq y_2 \geq \dots \geq y_n;$$

$$x_1 + x_2 + \dots + x_k \leq y_1 + y_2 + \dots + y_k \quad (1 \leq k \leq n-1).$$

The theory of majorization originated with the work of Schur (1923) and Hardy, Littlewood, and Pólya (1929), and is now firmly established; see Marshall and Olkin (1979). The following theorem was essentially proved by Schur (1923). A continuous analogue was proved by Hardy, Littlewood, and Pólya (1929), and an important generalization provided by Karamata (1932).

THEOREM 2. *Let ϕ be a real-valued convex function defined on an interval $I \subset \mathbb{R}$. If $(x_1, \dots, x_n) \prec (y_1, \dots, y_n)$ on I^n , then*

$$\sum_{i=1}^n \phi(x_i) \leq \sum_{i=1}^n \phi(y_i). \quad (3.1)$$

If, in addition, ϕ is strictly convex on I , then equality in (3.1) occurs if and only if $x_i = y_i$ ($i = 1, \dots, n$).

Proof. See Marshall and Olkin (1979, Propositions 3.C.1 and 3.C.1.a.). ■

An important application of Karamata's inequality is given in Theorem 3.

THEOREM 3. Let $A = (a_{ij})$ be a real symmetric $n \times n$ matrix with eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$. Then for any convex function $\phi: \mathbb{R} \rightarrow \mathbb{R}$ we have

$$\sum_{i=1}^n \phi(a_{ii}) \leq \sum_{i=1}^n \phi(\lambda_i). \quad (3.2)$$

Moreover, if ϕ is strictly convex, then equality in (3.2) occurs if and only if A is diagonal.

Proof. Without loss of generality we may assume that

$$\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n, \quad a_{11} \geq a_{22} \geq \dots \geq a_{nn}.$$

It is well known that (a_{11}, \dots, a_{nn}) is majorized by $(\lambda_1, \dots, \lambda_n)$; see Schur (1923) or Marshall and Olkin (1979, Theorem 9.B.1). Thus (3.2) follows from Theorem 2. For strictly convex ϕ , Theorem 2 implies that equality in (3.2) holds if and only if $a_{ii} = \lambda_i$ for $i = 1, \dots, n$, and, by Theorem 1, this is the case if and only if A is diagonal. ■

As a corollary of Theorem 3, let us establish the following useful inequalities concerning positive semidefinite matrices, which we shall use in proving Theorem 5.

THEOREM 4. Let $A = (a_{ij})$ be a positive semidefinite $n \times n$ matrix. Then

$$\operatorname{tr} A^p \geq \sum_{i=1}^n a_{ii}^p \quad (p > 1)$$

and

$$\operatorname{tr} A^p \leq \sum_{i=1}^n a_{ii}^p \quad (0 < p < 1)$$

with equality if and only if A is diagonal.

Proof. Let $p > 1$, and define $\phi(x) = x^p$ ($x \geq 0$). The function ϕ is strictly convex. Hence Theorem 3 implies that

$$\operatorname{tr} A^p = \sum_{i=1}^n \lambda_i^p(A) = \sum_{i=1}^n \phi(\lambda_i(A)) \geq \sum_{i=1}^n \phi(a_{ii}) = \sum_{i=1}^n a_{ii}^p$$

with equality if and only if A is diagonal. Next, let $0 < p < 1$. Then $\psi(x) \equiv -x^p$ ($x \geq 0$) is strictly convex, and the second result follows in the same way. ■

4. QUASILINEAR REPRESENTATION OF $(\operatorname{tr} A^p)^{1/p}$

We now have all the ingredients to prove the matrix analogue of Lemma 1.

THEOREM 5. *Let $p > 1$, $q = p/(p-1)$, and let $A \neq 0$ be a positive semidefinite $n \times n$ matrix. Then*

$$\operatorname{tr} AX \leq (\operatorname{tr} A^p)^{1/p} \quad (4.1)$$

for every positive semidefinite $n \times n$ matrix X satisfying $\operatorname{tr} X^q = 1$. Equality in (4.1) occurs if and only if

$$X^q = \frac{1}{\operatorname{tr} A^p} A^p. \quad (4.2)$$

REMARK. If we let R be the region

$$R = \{ X : X \in \mathbb{R}^{n \times n}, X \text{ positive semidefinite, } \operatorname{tr} X^q = 1 \},$$

then we can state Theorem 5 equivalently as

$$\max_R \operatorname{tr} AX = (\operatorname{tr} A^p)^{1/p} \quad (4.3)$$

for every positive semidefinite $n \times n$ matrix A . Thus we can express $(\operatorname{tr} A^p)^{1/p}$ (a *nonlinear* function of A) as an envelope of *linear* functions of A . In other

words, we obtain a quasilinear representation of $(\operatorname{tr} A^p)^{1/p}$.

Proof. Let X be an arbitrary positive semidefinite $n \times n$ matrix satisfying $\operatorname{tr} X^q = 1$. Let S be an orthogonal matrix such that $S'XS = \Lambda$, where Λ is diagonal and has the eigenvalues of X as its diagonal elements. Define $B = (b_{ij}) = S'AS$. Then

$$\operatorname{tr} AX = \operatorname{tr} B\Lambda = \sum b_{ii}\lambda_i$$

and

$$\operatorname{tr} X^q = \operatorname{tr} \Lambda^q = \sum \lambda_i^q.$$

Hence, by Lemma 1,

$$\operatorname{tr} AX = \sum b_{ii}\lambda_i \leq \left(\sum b_{ii}^p \right)^{1/p}. \quad (4.4)$$

Since A is positive semidefinite, so is B , and Theorem 4 thus implies that

$$\sum b_{ii}^p \leq \operatorname{tr} B^p. \quad (4.5)$$

Combining (4.4) and (4.5) we obtain

$$\operatorname{tr} AX \leq (\operatorname{tr} B^p)^{1/p} = (\operatorname{tr} A^p)^{1/p}.$$

Equality in (4.4) occurs if and only if

$$\lambda_i^q = \frac{b_{ii}^p}{\sum b_{ii}^p} \quad (i = 1, \dots, n),$$

and equality in (4.5) if and only if B is diagonal. Hence, equality in (4.1) occurs if and only if

$$\Lambda^q = \frac{B^p}{\operatorname{tr} B^p},$$

which is equivalent to (4.2). This concludes the proof. ■

5. MATRIX ANALOGUES OF THE INEQUALITIES OF HÖLDER AND MINKOWSKI

An immediate consequence of Theorem 5 is the matrix analogue of Hölder's inequality (1.2).

THEOREM 6. *For any two positive semidefinite matrices A and B of the same order, $A \neq 0$, $B \neq 0$, and $0 < \alpha < 1$, we have*

$$\operatorname{tr} A^\alpha B^{1-\alpha} \leq (\operatorname{tr} A)^\alpha (\operatorname{tr} B)^{1-\alpha}, \quad (5.1)$$

with equality if and only if $B = \mu A$ for some scalar $\mu > 0$.

Proof. Let $p = 1/\alpha$, $q = 1/(1 - \alpha)$, and assume $B \neq 0$. Now define

$$X = \frac{B^{1/q}}{(\operatorname{tr} B)^{1/q}}.$$

Then $\operatorname{tr} X^q = 1$, and hence Theorem 5 applied to $A^{1/p}$ yields

$$\operatorname{tr} A^{1/p} B^{1/q} \leq (\operatorname{tr} A)^{1/p} (\operatorname{tr} B)^{1/q},$$

which is (5.1). According to Theorem 5, equality in (5.1) can only occur if $X^q = (1/\operatorname{tr} A)A$, that is, if $B = \mu A$ for some $\mu > 0$. ■

Another consequence of Theorem 5, more specifically of its quasilinear representation (4.3), is the matrix version of Minkowski's inequality (1.4).

THEOREM 7. *For any two positive semidefinite matrices A and B of the same order ($A \neq 0$, $B \neq 0$), and $p > 1$, we have*

$$[\operatorname{tr} (A + B)^p]^{1/p} \leq (\operatorname{tr} A^p)^{1/p} + (\operatorname{tr} B^p)^{1/p}. \quad (5.2)$$

with equality if and only if $A = \mu B$ for some $\mu > 0$.

Proof. Let $p > 1$, $q = p/(p - 1)$, and let R be the region

$$R = \{ X : X \in \mathbb{R}^{n \times n}, X \text{ positive semidefinite, } \operatorname{tr} X^q = 1 \}.$$

Using the quasilinear representation (4.3) of $(\operatorname{tr} A^p)^{1/p}$,

$$\max_R \operatorname{tr} AX = (\operatorname{tr} A^p)^{1/p},$$

we obtain

$$\begin{aligned} [\operatorname{tr}(A+B)^p]^{1/p} &= \max_R \operatorname{tr}(A+B)X \\ &\leq \max_R \operatorname{tr} AX + \max_R \operatorname{tr} BX \\ &= (\operatorname{tr} A^p)^{1/p} + (\operatorname{tr} B^p)^{1/p}. \end{aligned} \quad (5.3)$$

Equality in (5.3) can only occur if the same X maximizes $\operatorname{tr} AX$, $\operatorname{tr} BX$, and $\operatorname{tr}(A+B)X$, which implies, by Theorem 5, that A^p , B^p , and $(A+B)^p$ are proportional, and hence that A and B are proportional. ■

I am very grateful to Ken Binmore for his constructive remarks.

REFERENCES

- Beckenbach, E. F. and Bellman, R. 1961. *Inequalities*, Springer, Berlin.
- Hardy, G. H., Littlewood, J. E., and Pólya, G. 1929. Some simple inequalities satisfied by convex functions, *Messenger Math.* 58:145–152.
- Hardy, G. H., Littlewood, J. E., and Pólya, G. 1952. *Inequalities* (2nd ed.), Cambridge U.P., Cambridge.
- Karamata, J. 1932. Sur une inégalité relative aux fonctions convexes, *Publ. Math. Univ. Belgrade* 1:145–148.
- Marshall, A. W. and Olkin, I. 1979. *Inequalities: Theory of Majorization and Its Applications*, Academic, New York.
- Schur, I. 1923. Ueber eine Klasse von Mittelbildungen mit Anwendungen auf die Determinantentheorie, *Sitzungsber. Berlin. Math. Ges.* 22:9–20.

Received 23 December 1985; revised 17 November 1986